

ENVIRONMENTALLY FRIENDLY NUTRIENT SUPPLY OF SPRING BARLEY

¹PÉTER JAKAB, ²PIROSKA NAGY JAKABNÉ

¹ University of Szeged Faculty of Agriculture,
H-6800 Hódmezővásárhely, Andrásy u. 15., Hungary
² College of Szolnok, H-5000 Szolnok, Tiszaligeti St. 14., Hungary
jakabpeter@mgk.u-szeged.hu

ABSTRACT

We examined the effect of fertilization and soil bacterium preparation on the yield of spring barley varieties in 2011. The experiment was set in three repetitions, random blocks. The soil of the experiment was calcareous meadow chernozem. During the experiment we applied seven different fertilizer bacteria treatments, which we supplemented with a control plot. The year 2011 was unfavourable for spring barley production. In 2011 the amount of precipitation in the vegetation period of spring barley was lower than the average of 30 by 39.7 mm. As a result, relatively low yields formed. The control plots had the lowest average yield, 2.2-2.42 t/ha. We can claim that improving the nutrient supply the crop results were higher. Without the bacteria treatment the crop results were between 3.12 and 4.42 t/ha. The most favourable results occurred in case of N 80 kg/ha+PK and N 120kg/ha+PK treatments. Under the influence of soil bacteria treatments the average yields were better, which obviously proves the nutrient transformation ability of the bacteria. With bacteria treatments the result was 3.37 –4.5 t/ha, so owing to this we registered 0.1 – 0.44 t/ha extra yield.

Keywords: spring barley, nutrient supply, yield, fertilization, soil bacterium preparation

INTRODUCTION

The ecological conditions of Hungary are favourable for cereals production. Spring barley is a plant produced for the brewing industry and for animal feeding purposes. Unlike other spring crops it shows unusual temperature needs and more moderate water requirements (ca. 225-250 mm in the vegetation period), but it is sensitive to the distribution of precipitation (KISMÁNYOKI, 2005).

The yield of arable crops is influenced by many factors; among them the nutrients play a determinative role in intensive field crop production (PEPÓ, 2007; PETRÓCZI, 2008).

To provide the required amount of supplementary nutrients, it is necessary to determine the suitable NPK nutrient rate for the plants. Nitrogen can be absorbed only in the presence of the necessary phosphorus and potassium supplies, and large dose of nitrogen alone induces crop depression (SÁRVÁRI, 2006).

The short-term breeding spring barley has poorly developed roots, and therefore it needs a lot of nutrients it can take up easily. To provide the high yield and good quality it is very important to ensure the optimal amount and ratio of nutrients, suitable for production purposes. The nutrient supply, the ratio and the quantities of nutrients (NPK) have fundamental influence on the yield and quality (RADICS, 1994).

The microbial vaccines mainly provide nutrients supply and plant growth-promoting hormones as well as breaking down raw materials. In addition, they have an activity against soil and seed-derived microbes (ÁRVAY, 2004).

One factor, which prevents the use of vaccines, is that the effect of microbicide vaccines is less spectacular, especially compared with the results of the use of fertilizers. In many cases, the products affect only the following six crops, e.g. improve the soil structure, or the residual effects of nutrients (BIRÓ, 2004).

The use of soil bacteria preparations has several advantages. Some of the bacteria they contain bind the nitrogen in the air, while others explore phosphorus and potassium contents that plants cannot take up. This results in significant fertilizer-cost savings. Other

bacteria reduce agents in the soil, thereby increasing the resistance of plants and reduce the number of fungicide preventions. As a result, less pesticides and fungicides will be necessary. They improve the soil structure, thereby improving physical and chemical properties of the soil. The lighter soils can take the precipitation better, reducing the risk of developing inland waters. On the other hand, a higher moisture content of the soil in this area would reduce drought sensitivity. The soils treated with soil bacteria preparation, due to its improving soil structure, have lower resistance to the tillage tool, which reduces the amount of fuel used for tillage. They are capable of producing plant hormones and as a result they contribute to the improved drought tolerance of plants as well. The above advantages of the soil bacterial products are justified in any field crop production. (AGRO.BIO HUNGARY KFT, 2013).

MATERIAL AND METHOD

Soil properties of the experimental field

We set the experiment on the area next to the Mezőtúr Industrial Park in 2011. The soil was calcareous meadow chernozem, the reaction of which was nearly neutral (pH_{KCL} 6.72). Before setting the experiment the soil analysis data showed that it had proper nitrogen, poor phosphor and good potassium content. The soil was hard to cultivate, with slow transformation of phosphor and potassium (*Table 1.*).

Table 1. Main properties of the experimental field area

pH (H₂O)	CaCO₃	P₂O₅ (mg/kg)	K₂O (mg/kg)	Humus (%)	Soil plasticity value (K_A)
6.72	0.7	64	433	3.07	57

Weather in the experimental years

The year 2011 was unfavourable for spring barley production. In 2011 the amount of precipitation in the vegetation period of spring barley was lower than the average of 30 by 39,7 mm. Particularly April and May were extremely dry when only 11.2 mm and 33.3 mm rain fell. Totally, we can say, the deficient precipitation had a negative effect on the development of spring barley, which resulted in low yields in 2011 (*Table 2.*).

Table 2. The distribution of precipitation in the vegetative period of spring barley in 2011

Month	Rainfall (mm)	Average rainfall (mm)	Difference (mm)
March	25.8	33	-7.2
April	11.2	46	-34.8
May	33.3	56	-22.7
June	56.6	59	-2.4
July	77.4	50	27.4
Total amount of rainfall (mm)	204.3	244	-39.7

Main features of the agrotechnique applied

Our small-scaled plough experiment was set in three replications, organised as a random block in 2011. We applied seven different fertilizer bacteria treatments, which we supplemented with a control plot. The amount of nitrogen was applied in autumn and spring in 50-50 %; the total amount of phosphorus and potassium was applied in autumn in one dosage. The soil bacterium preparation was applied in spring. The fore-crop was maize. Fall tillage involved deep ploughing at 28-32 cm depth in the experimental years. The spring barley varieties in the experiment were Pasadena, KH Lédi, KH Szofi, and Mauritia.

We processed the obtained data by single factor variant analysis (SVÁB, 1981).

RESULTS

Without any fertilizers the yield of the examined varieties was ranging between 2.2-2.42 t/ha. The KH Szofi had the best ability to absorb and utilize nutrients. It obtained a yield of 2.42 t/ha without any fertilizers. The yield of varieties was ranging between 3.12-4.42 t/ha in the consequence of fertilization. By assessing the varieties we can conclude, that the KH Szofi reached the highest yield with 4.42 t/ha in N 120+PK treatment. The economical and reliable yield increase was achieved by N 40+PK and N 80+PK treatments. Under the influence of soil bacteria treatments the average yields were better, which obviously proves the nutrient transformation ability of the bacteria. With bacteria treatments the result was 3.37 –4.5 t/ha, so owing to this we registered 0.1 – 0.44 t/ha extra yield (*Figure 1.*)




Figure 1. The yield of spring barley varieties on different nutrient levels

CONCLUSIONS

Under the influence of soil bacteria treatments the yield was better, which obviously proves the nutrient transformation ability of the bacteria. Proper conclusion can be drawn only from the results of several years this making further examinations is essential.

We carried out our research from the conviction that the results obtained should be utilized on the geographical areas with similar ecological character; furthermore, newer information shall be obtained that help in the elaboration and specification of technologies for each variety, respectively.

ACKNOWLEDGEMENTS

We thank the College of Szolnok for financial support of this experiment.

REFERENCES

- AGRO.BIO HUNGARY KFT (2013): Termékek. <http://agrobio.hu/?base=products>, 2013.02.21.
- ÁRVAY, GY. (2004): Baktériumtrágyák. Mezőgazdasági tanácsok 13. évf. pp. 10-12.
- BÍRÓ B. (2004): Vegyük, vagy ne vegyük? A baktériumtrágyák haszna és alkalmazási szempontjai. Mezőgazdasági tanácsok 13. évf. pp. 12-15.
- KISMÁNYOKY T. (2005): Őszi és tavaszi árpa. In: Antal J (ed.) Növénytermesztés 1. A növénytermesztés alapjai. Gabonafélék. Mezőgazda Kiadó, Budapest. pp. 244-276.p.
- PEPÓ, P. (2007): The role of fertilization and genotype in sustainable winter wheat (*Triticum aestivum* L.) production. Cereal Research Communications, Volume 35. Number 2. pp. 917-920.
- PETRÓCZI, I. M. (2008): Termeljünk több búzát! – Hogyan? Agroforum. 19. évf. 7/M. sz. pp. 2.
- RADICS L. (1994): Szántóföldi növénytermesztés. Egyetemi jegyzet. Budapest. pp. 63-64.
- SÁRVÁRI, M. (2006): A vetésforgó és a tápanyagellátás szerepe az őszi búza termesztésében. In: Búzavertikum aktuális kérdései. Szaktanácsadási füzetek 2. Debrecen. pp. 64-72
- SVÁB, J. (1981): Biometria módszerek a kutatásban. Mezőgazdasági Kiadó. Budapest. 557. p.